

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/62, C07K 14/71, 14/715, 16/28, G01N 33/68		A1	(11) International Publication Number: WO 95/07984
			(43) International Publication Date: 23 March 1995 (23.03.95)
(21) International Application Number: PCT/US94/10457		(81) Designated States: AT, AU, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KR, LT, LU, LV, NL, NO, NZ, RU, SE, SI, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 15 September 1994 (15.09.94)			
(30) Priority Data: 08/122,796 16 September 1993 (16.09.93) US		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant: INDIANA UNIVERSITY FOUNDATION [US/US]; Showalter House, P.O. Box 500, Bloomington, IN 47402 (US).			
(72) Inventor: KWON, Byoung, S.; Indiana School of Medicine, Medical Science, Room 255, 63 Barnhill Drive, Indianapolis, IN 46223 (US).			
(74) Agent: MICHAELS, Christopher, A.; Barnard, Brown & Michaels, Suite 220, 306 E. State Street, Ithaca, NY 14850 (US).			

(54) Title: HUMAN RECEPTOR H4-1BB

(57) Abstract

The human receptor H4-1BB has been isolated, sequenced and disclosed herein. The cDNA of the human receptor H4-1BB is about 65 % homologous to the mouse cDNA 4-1BB and was isolated by using probes derived from cDNA 4-1BB. A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB was developed. It comprises the extracellular portion of the receptor protein H4-1BB and a detection protein (alkaline phosphatase) bound to the portion of the receptor protein H4-1BB. B-cells that have expressed a ligand to receptor protein H4-1BB can be treated with cells that have expressed receptor protein H4-1BB and B-cell proliferation may be induced. The use of H4-1BB to block H4-1BB ligand binding has practical application in the suppression of the immune system during organ transplantation. A monoclonal antibody against H4-1BB can be used to enhance T-cell proliferation by treating T-cells that have expressed receptor protein H4-1BB with the anti H4-1BB monoclonal antibody. Tumors transfected with H4-1BBL may be capable of delivering antigen-specific signals as well as the co-stimulatory signals and can be killed by human cytotoxic T lymphocytes.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

HUMAN RECEPTOR H4-1BB

5

This application is a continuation-in-part of co-pending application Serial No. 08/012,269 filed February 1, 1993, which is a continuation-in-part of co-pending application Serial No. 07/922,996 filed July 30, 1992, 10 which is a continuation-in-part of copending application Serial No. 07/267,577 filed November 7, 1988.

The subject matter described herein was in part a subject invention of NIH Grants Nos. IR23AI23058-03, R01 AI28175 and P60 KD20542 of which the present inventor was 15 the Principal Investigator and either the Donald Guthrie Foundation for Medical Research Inc. of Guthrie Square, Sayre, Pennsylvania 18849-1669 or Indiana University School of Medicine of Indianapolis, Indiana 46202, was the Grantee.

20

FIELD OF THE PRESENT INVENTION

The present invention relates to a previously unknown 25 human receptor protein, H4-1BB, which was isolated and identified based upon work with a homologous murine (mouse) receptor protein, 4-1BB, which was isolated and identified by specific expression of the T cell genes by the present inventor.

30

BACKGROUND OF THE PRESENT INVENTION

The immune system of humans and other species requires that white blood cells be made in the bone marrow, which 35 white blood cells include phagocytes, lymphocytes and B cells. As presently understood, the phagocytes include macrophage cells which scavenge unwanted materials such as

virus protein from the system. The lymphocytes include helper T cells and killer T cells and B cells as well as other cells, including those categorized as suppressor T cells. The B cells produce the antibodies. The killer T
5 cells physically pierce the cell and the helper T cells facilitate the whole process. In any event, the immune process is facilitated by lymphokines.

Lymphokines are the proteins by which the immune cells communicate with each other. Scientists produce them in
10 sufficient quantities for therapeutic use against immunologic diseases. There are many known lymphokine proteins and they include the interferons, interleukin-1,2,3,4,5,6,7, colony-stimulating factors, lymphotoxin, tumor necrosis factor and erythropoietin, as well as
15 others.

Interleukin 1, secreted from macrophages activate the helper T cells and raise the body temperature causing fever which enhances the activity of the immune cells. The activated helper T Cells produce Interleukin 2 and
20 Interleukin 2 stimulates the helper and killer T cells to grow and divide. The helper T cells also produce another lymphokine, B cell growth factor (BCGF), which causes B cells to multiply. As the number of B cells increases, the helper T cells produce another lymphokine known as the B
25 cell differentiating factor (BCDF), which instructs some of the B cells to stop replicating and start producing antibodies. T cells also produce a lymphokine, gamma interferon (IF), which has multiple effects like Interleukin 2. Interferon helps activate killer T cells,
30 enabling them to attack the invading organisms. Like BCGF, interferon increases the ability of the B cells to produce antibodies. Interferon also affects the macrophages to keep them at the site of the infection and help the macrophages to digest the cells they have engulfed.
35 Gathering momentum with each kind of lymphokine signal between the macrophages and the T cells, the lymphokines amplify the immune system response and the virus protein or

other foreign matter on the infected cells is overwhelmed. There are many other lymphokines, maybe a hundred or more, which participate in the immune process. Many lymphokines are known and many are not.

5 Lymphokines are sometimes called intercellular peptide signals. Among scientists there is widespread use of cloned cell lines as lymphokine producers and the isolation of lymphokine mRNA has become a common technique. The mouse
10 receptor protein, 4-1BB, was isolated and identified based on specific expression of the T cell genes using a technique identified by the present inventor in a publication (Proc. Natl. Acad. Sci. USA. 84, 2896-2900, May 1987, Immunology). The protocol reported in this
15 publication can be used by scientists to detect virtually all of the lymphokines. The method is designed to detect virtually all mRNA expressed differentially and the mRNA sequences of the immune cells are expressed differentially (as they relate to the T cells and the killer T cells) even though the level of expression is low and the quantity of
20 the secreted lymphokine protein is low. The present inventor believes that the analysis described in the above identified publication can reveal biologically important molecules such as lymphokines because there are many indications that biologically important or active molecules
25 are coded by the most scarce messages. An example is a transforming growth factor (TGF) which is present as only one of a million clones.

Most T cell factors have been classically identified by recognizing biologic activities in assays, purifying the
30 protein information. An alternative approach is to isolate putative T cell genes based upon specific expression and then demonstrate the function of the unknown molecule. Using the aforesaid modified differential screening procedure, the present inventor cloned a series of T cell
35 subset-specific cDNAs from cloned helper T (HTL) L2 and cloned cytolytic T lymphocyte (CTL) L3.

A series of T-cell subset-specific cDNAs were isolated from cloned murine T-cells by employing a modified differential screening procedure. The nucleotide sequence and expression properties of some of the cDNA species have been reported. One of the genes not previously characterized, that encodes mouse receptor protein 4-1BB, was studied further. These studies have led to the isolation of the human homologue to 4-1BB, H4-1BB.

10

SUMMARY OF THE PRESENT INVENTION

The present invention includes the human receptor protein H4-1BB and the cDNA gene encoding for human receptor protein H4-1BB. The nucleotide sequence of the isolated cDNA is disclosed herein along with the deduced amino acid sequence. The cDNA gene identified as pH4-1BB was deposited at the Agricultural Research Service Culture Collection and assigned the accession number: NRRL B21131

The cDNA, and fragments and derivatives thereof, can be used as a probe to isolate DNA sequences encoding for proteins similar to the receptor protein encoded by the cDNA. The cDNA of the human receptor H4-1BB is about 65% homologous to the mouse cDNA 4-1BB and was isolated by using probes derived from cDNA 4-1BB. The cDNA gene identified as p4-1BB was deposited at the American Type Culture Collection at 12301 Parklawn Drive, Rockville, Maryland 20852 under ATCC No.: 67825.

The human receptor protein H4-1BB can be produced by:

- 1) inserting the cDNA of H4-1BB into an appropriate expression vector, 2) transfecting the expression vector into an appropriate transfection host, c) growing the transfected hosts in appropriate culture media and d) purifying the receptor protein from the culture media. The protein and fragments and derivatives can be used: 1) as a probe to isolate ligands to human receptor protein H4-

1BB, 2) to stimulate proliferation of B-cells expressing H4-1BB ligands, or 3) to block H4-1BB ligand binding.

B-cell proliferation can be induced by treating B-cells that have expressed a ligand to receptor protein H4-1BB with cells that have expressed receptor protein H4-1BB. The use of H4-1BB to block H4-1BB ligand binding has practical application in the suppression of the immune system during organ transplantation. A similar costimulatory immune system pathway is being analyzed for this type of application. See "Mounting a Targeted Strike on Unwanted Immune Responses", Jon Cohen, Science, Vol. 257, 8-7-92; "Long Term Survival of Xenogeneic Pancreatic Islet Grafts Induced by CTLA4Ig", Lenschow et al, Science Vol. 257, 7-8-92; and "Immunosuppression in Vivo by a Soluble Form of the CTLA-4 T Cell Activation Molecule", Linsley et al, Science Vol. 257 7-8-92.

A monoclonal antibody against H4-1BB can be used to enhance T-cell proliferation by treating T-cells that have expressed receptor protein H4-1BB with the anti H4-1BB monoclonal antibody. Some tumors are potentially immunogenic but do not stimulate an effective anti-immune response in vivo. Tumors may be capable of delivering antigen-specific signals to T cells, but may not deliver the co-stimulatory signals necessary for full activation of T cells. Expression of the co-stimulatory ligand B7 on of melanoma cells was found to induce the rejection of a murine melanoma in vivo. ("Tumor Rejection After Direct Co-Stimulation of CD8⁺ T Cells by B7-Transfected Melanoma Cells", Sarah E. Townsend and James P. Allison, Science Vol. 259, 1-5-93.) A monoclonal antibody against H4-1BB may be capable of the same effect as it is now known to induce T cell proliferation and activation.

A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB was developed. It comprises the extracellular portion of the receptor protein H4-1BB and a detection protein (alkaline phosphatase) bound to the portion of the receptor protein H4-1BB. The portion

of the receptor protein H4-1BB binds to the cell membrane ligands and binding can be detected by relative activity assays for the detection protein. The fusion protein is placed in the presence of a cell suspected to express the
5 receptor protein H4-1BB. Then the cell is washed of any fusion protein not bound to the cell membrane ligands. Once the washed cells are placed in the presence of a substrate for the detection protein and the relative activity of the detection protein can be measured.

10 The primary object of the present invention is the identification of the new human receptor, H4-1BB as identified herein by its sequence.

Another object of the present invention is to teach a fusion protein comprising the extracellular portion of H4-
15 1BB and a detection protein.

Still another object of the present invention is to teach methods of using the cDNA H4-1BB, the receptor protein H4-1BB, the monoclonal antibody and the legend for H4-1BB.

20

BRIEF DESCRIPTIONS OF THE FIGURES

Figure 1 shows the sequence for the cDNA of mouse
25 receptor protein 4-1BB and the regions used as PCR primers to obtain the human homologue H4-1BB.

Figures 2a and 2b show the nucleotide sequence and the deduced amino acid sequence of human receptor H4-1BB respectively.

30 Figures 3a and 3b illustrate the molecules involved in T-cell activation.

Figures 4a, 4b, and 4c illustrate a normal T-cell activation pathway.

Figures 5a, 5b, and 5c illustrate CTLA4-lg alone, 4-
35 1BB/AP and CTLA4-lg together and 4-1BB/AP alone respectively being used to block steps in the T-cell activation pathway.

DETAILED DESCRIPTION

In the following detailed description references are made to known procedures and studies, as well as published work of the applicant. These publications are incorporated herein by reference for clarity and listed in an appendix included at the end of this detailed description.

10 Isolation and characterization of mouse receptor 4-1BB

Figure 1 shows the nucleotide sequence and the deduced amino acid sequence of the mouse receptor 4-1BB. The nucleotides of the message strand are numbered in the 5' to 3' direction and numbers are shown on both sides of the sequence. Nucleotide residue 1 is the A of the initiation codon ATG, and the nucleotides on the 5' side of residue 1 are indicated by negative numbers. The predicted amino acid sequence is shown below the nucleotide sequence. Putative signal peptide is underlined. Stop codon is indicated by (---). Cysteine residues are highlighted by the dots. An unusual feature of 4-1BB sequence is that there is a potential polyadenylation signal of AATAAA at nucleotides 1158-1163 (Fig. 1 boxed). It was believed that this signal was functional because this gene produces at least two different sizes of mRNA.

The transcript of 4-1BB was inducible by concanavalin A in mouse splenocytes, T-cell clones, and hybridomas. The expression of 4-1BB transcripts was inhibited by cyclosporin A. The 4-1BB mRNA was inducible by antigen receptor stimulation but was not inducible by Il-2 stimulation in the cloned T-cells (1). The 4-1BB cDNA encodes a peptide of 256 amino acids containing a putative leader sequence, a potential membrane anchor segment, and other features of known receptor proteins. Therefore, the expression pattern of 4-1BB resembles those of lymphokine mRNAs while the sequence appeared consistent with those of receptor proteins.

The major species of 4-1BB on the cell surface appears to be a 55-kDa dimer. 4-1BB also appears to exist as a 30-kDa monomer and possibly as a 110-kDa tetramer. Since these 4-1BB species were immunoprecipitated from a homogenous population of cells (T cell clone F1), all forms potentially co-exist on each cell. A comparison of peptide digests from the 4-1BB monomer and dimer will be needed to determine whether 4-1BB exists as a homodimer on the cell surface. A variety of cell surface receptors such as the insulin receptor (2), the B cell surface immunoglobulin receptor (3), the T cell Ag receptor (4), the CD28 costimulatory receptor (5), and the CD27 T cell antigen (6) are composed of disulfide-bonded subunits. Receptor dimerization may be required for ligand binding and subsequent biochemical signaling.

4-1BB is not expressed on resting T cells but is inducible by activators which deliver a complete growth stimulus to the T cell. The combination of PMA and ionomycin is capable of mimicing those signals required for T cell proliferation. Although PMA or ionomycin alone induced 4-1BB mRNA, the combination of PMA and ionomycin resulted in optimal 4-1BB expression. Furthermore, the expression of 4-1BB was not transient. When purified splenic T cells were stimulated with immobilized anti-CD3, 4-1BB mRNA was expressed and this expression was maintained for up to 96 hrs poststimulation. Cell cycle analysis will be required to confirm that 4-1BB is expressed throughout cell cycle progression.

4-1BB is structurally related to members of the nerve growth factor receptor super-family. Although these receptors possess structurally similar ligand-binding properties (cysteine-rich regions), the cytoplasmic domains of these proteins are nonconserved which could allow for diversity in transmembrane signaling. Some members of this family are involved in the T or B cell activation process. There are in vitro functional data on the OX-40, CD40 and CD27 antigens. Antibodies against the OX-40 augment the T

cell response in a mixed lymphocyte reaction (7) and antibodies against CD40 enhance B-cell proliferation in the presence of a coactivator, such as PMA or CD20 antibodies, and synergize with IL-4 in vitro to induce B-cell differentiation and to generate long-term normal B cell lines (8). One monoclonal antibody, anti-1A4, which recognizes an epitope on the CD27 molecule inhibited calcium mobilization, IL-2 secretion, helper T cell function, and T cell proliferation. On the other hand, CLB-CD27/1, another anti-CD27 mAb enhanced proliferation of human T cells stimulated with PHA or anti-CD3 mAb (6). These results indicate that the CD27 molecule plays an important role in T cell activation. Except for TNFRs, NCAM and CD40, the ligands or cell surface molecules to which the members of the superfamily bind are not yet identified. Identification and characterization of the ligands to which the receptors bind will be helpful in better defining the physiologic role of 4-1BB.

To ascertain whether cell surface 4-1BB could contribute to T cell activation, the anti-4-1BB 53A2 was used as an antagonist to 4-1BB. These data suggested that 4-1BB does in fact have the potential to function as an accessory signaling molecule during T cell activation and proliferation. The addition of soluble 53A2 to purified splenic T cells stimulated with immobilized anti-CD3 resulted in an amplification of ³H thymidine incorporation compared to T cells stimulated with anti-CD3 alone. This pattern of enhancement ranged from 2- to 10- fold in three independent experiments.

In the original two signal model of Bretscher and Cohn, they proposed that signal 1, the occupancy of the T cell antigen receptor (TCR), resulted in inactivation of the T cell in the absence of signal 2, which is provided by accessory cells. This has since been confirmed by a variety of studies (9). The identification of the accessory cell CD28 as a potent costimulatory receptor on T cells was a significant contribution in beginning to characterize the

accessory signal(s) required for optimal T cell proliferation (10). It is possible that other cell surface molecules may contribute to these costimulatory activation requirements (11).

5 The biochemical signals delivered through 4-1BB are not completely known. One possibility considered was the observation that 4-1BB contains a putative p56^{lck} tyrosine kinase binding domain in its cytoplasmic tail. It was later determined that p56^{lck} tyrosinase kinase binds to
10 4-1BB. It will also be worthwhile to determine if 4-1BB-mediated signaling can regulate genes such as IL-2 and IL-2 receptor, whose expression is required for T cell activation and subsequent proliferation.

Although the precise functions of members of the Nerve
15 Growth Factor Receptor (NGFR) family appear to be diverse, an emerging theme is one in which these molecules may contribute in various ways to a maintenance of responsiveness or viability of the particular cell type in which they are expressed. For instance, NGF is absolutely
20 required for viability of neurons in vitro and in vivo (12). The crosslinking of CD40 by soluble antiCD40 monoclonal antibody blocks germinal center centrocytes from undergoing apoptosis in vitro (13). Signals delivered through CD40 may also aid in maintenance of responsiveness
25 to differentiation factors. The ligation of CD40 with anti-CD40 F(ab')₂ fragments in the presence of IL-4 induced large increases IgE synthesis (14). Also, anti-CD40 activated naive B cells treated with IL-10 and transforming growth factor- β became committed to IgA secretion (15).

30 In addition to sharing the molecular characteristics with the NGFR superfamily, it was noted that the 4-1BB contained a putative zinc finger structure of the yeast eIF-2 β protein (16). 4-1BB also shares a conserved region with the *sina* seven in absentia of *Drosophila*, which is
35 required for correct photoreceptor cell development (17). That particular region is also similar to the protein

product of the DGl7 gene of *Dictyostelium*, whose expression is specifically induced during aggregation by cAMP (18).

This region forms the pattern of C-X₂-C-X₂-C-X₃-H-X₃-C-X-C; and the cysteines and histidine are conserved in a similar space in 4-1BB, *sina*, and DGl7 proteins. Ten of 24 amino acids between the 4-1BB and *sina* proteins are identical, and 3 of 24, are conservative substitutes. The conserved pattern suggests that these amino acids are functionally important. The *sina* protein is localized in the nucleus, suggesting that it has a regulatory function in cells. The fact that the amino acid sequence of 4-1BB contains features like a zinc finger motif, a nuclear protein, and a receptor domain suggests that 4-1BB may play diverse roles during cellular proliferation and differentiation.

4-1BB may represent another cell-surface molecule involved in T cell-APC interactions. The 4-1BB-AP fusion protein specifically bound to mature B-cell lines, anti- μ -activated primary B cells, and mature macrophage-cell lines. 4-1BB-AP bound at low or insignificant levels to immature B- and macrophage-cell lines, T-cell clones, T-cell lines, primary culture T cells, and various nonlymphoid-cell lines. Since 4-1BB-AP binds to mature B cells and macrophages, it is possible that signals delivered upon 4-1BB binding may modulate APC functions in some way. This possibility remains to be explored.

Chalupny and colleagues (19) have proposed that 4-1BB Rg, a fusion protein consisting of the extracellular domain of 4-1BB and the Fc region of human IgG, bound to the extracellular matrix (ECM). The highest level of 4-1BB Rg binding was to human vitronectin. In data not shown, an ELISA was performed using 4-1BB-AP and human vitronectin (Yelios Pharmaceuticals/GIBCO-BRL, Grand Island, NY.) immobilized at 0.007 μ g-10 μ g per well on microtiter plates. No binding of 4-1BB-AP based on AP activity was observed. To rule out the possibility that 4-1BB-AP was binding to proteins extrinsically attached to the cell

surface (possible extracellular matrix components), B-cell lymphomas were washed in acid conditions prior to the binding assay. 4-1BB-AP still bound specifically to mature B-cell lymphomas. It is still to be determined whether a 4-1BB-ligand specifically expressed on B cells and macrophages exists, and whether 4-1BB-AP may bind to the ECM under particular binding conditions. It is possible that the ECM could facilitate the binding of 4-1BB to a specific cell-surface ligand.

10 B cells and helper T cells interact with each other through receptors on B cells binding to their specific counter-receptors on T cells. It is thought that this interaction results in a cascade of biochemical signaling relays between these two cell types (20). As this
15 interaction proceeds, these cells become committed to enter the S phase of the cell cycle. Initial interactions between TCR and CD4 on T cells, and processed antigen-MHC II on B cells, do not result in B cells capable of entering the cell cycle (21). However, studies from in vitro systems
20 suggest that once T-cells are stimulated, they express newly synthesized or modified cell-surface molecules capable of inducing B cells to enter the cell cycle (22, 23). This T-cell function is not antigen-specific or MHC-restricted (24). In addition, soluble factors are not
25 required for the activated Th induction of B-cell activation (25). Once B cells enter the cell cycle, IL-4 induces B cells to progress from G₁ to S phase. The ability of activated T cells or T-cell membranes to promote the entry of B cells into the cell cycle can be blocked by
30 either cycloheximide or cyclosporin A treatment (26, 27). These newly expressed membrane proteins appear to be "lymphokine-like" in their induction characteristics.

4-1BB has expression properties which meet the requirements of a B-cell costimulator. 4-1BB is inducible
35 by anti-CD3 or TCR-mediated T-cell stimulation, and its expression is sensitive to cyclosporin A as well as cycloheximide treatment (28). Interestingly,

paraformaldehyde-fixed SF21-4-1BB cells, synergized with anti- μ in inducing B-cell proliferation. The costimulation of splenic B cells by SF21-4-1BB occurred at optimal (10 μ g/ml) and suboptimal (1.0-0.1 μ g/ml) doses of anti- μ . The addition of SF21-4-1BB cells to resting B cells, did not result in significant B-cell proliferation. SF21-4-1BB cells did not synergize with TPA or ionomycin, or suboptimal concentrations of LPS in inducing B-cell proliferation.

Although the baculovirus system has been used to express large amounts of recombinant soluble proteins, this system may be utilized for the expression of recombinant cell-surface proteins. The baculovirus infection provides a convenient means to express uniformly high levels of recombinant protein on a per cell basis. It is noteworthy, that the addition of SF21 cells alone did not result in significant levels of costimulation. This can be a potential problem when using cos- or L- cell lines which can exhibit strong costimulator activity on their own.

Another member of the NGFR superfamily, CD40, is expressed on B cells and interacts with gp39, a molecule expressed on activated T cells. The cDNAs encoding the murine (29) and human (30) gp39 proteins have been cloned; this cell surface molecule is a type II membrane protein with homology to tumor necrosis factor. Noelle et al. (31) found that a CD40-inununoglobulin fusion protein, is capable of blocking T cell-induced B-cell proliferation and differentiation in a dose-dependent manner. Armitage et al. have isolated a cDNA for murine gp39 and showed that gp39 could induce B-cell proliferation in the absence of co-stimuli, and result in IgE production in the presence of IL-4-. Hollenbaugh et al. (32) have shown that COS cells transfected with human gp 39 can synergize with either TPA or anti-CD20 in inducing human B-cell proliferation and is able to stimulate B cells without a costimulator only at low levels. These data indicate that CD40 may be one of the

B-cell-surface molecules that transmit signals during physical contact with T cells.

Cell-surface receptors communicate with their external milieu by interacting either with soluble factors or other cell surface molecules expressed on neighboring cells. The role of biochemical signals delivered by cell-cell contact versus those delivered by soluble factors interacting with cell surface receptors is not clear. The NGFR superfamily is unusual for the TNFR I and II as well as the NGFR bind to more than one ligand. The TNFRs I and II both bind to TNF- α and TNF-R (33). The NGFR binds to NGF, brain-derived neurotrophic factor, and neurotrophin-3 (34).

In addition, one ligand may function as both a cell surface and soluble ligand. Recent evidence on the CD4-0 ligand, gp39, suggests that this ligand can exist as a membrane bound as well as a soluble ligand (35). It may be possible that 4-1BB is secreted and interacts with B cells in a soluble form as well as a membrane bound form. A member of the NGFR receptor family, CD27, which is expressed on T cells, is secreted in addition to being expressed on the cell surface (36). It is also possible that more than one ligand (soluble and cell surface) may bind to 4-1BB.

25 Isolation of the human homologue, H4-1BB

In order to isolate the human homologue (H4-1BB) of mouse 4-1BB two sets of polymerase chain reaction (PCR) primers were designed. To design the PCR primers, the amino acid sequence among the members of nerve growth factor receptor (NGFR) superfamily were compared because 4-1BB is a member of the superfamily (37). The amino acid sequences employed were mouse 4-1BB (38), human NGFR (39), human tumor necrosis factor receptors (33), human CD40 (40), and human CD27 (6). The areas of sequence conservation among the NGFR superfamily were chosen.

Forward primer I (H4-1BBFI) spans from amino acids 36 to 41 and forward primer II (H4-1BBFII) spans from amino

acids 52 to 58 of the mouse 4-1BB. Reverse primer I (H4-1BBRI) spans from amino acids 116 to 121 and reverse primer II (H4-1BBRII) spans from amino acids 122 to 128 of mouse 4-1BB. The regions used as PCR primers in mouse 4-1BB are indicated in Fig. 1.

The degenerative oligonucleotide sequence of each primer is as follows:

10	H4-1BBFI:	5' TTC TGT CGI AAA TAT AAT CC 3'
		T C A G C C
	H4-1BBFII:	5' TTC TCI TCI ATT GGI GGI CA 3'
		T G G C
		A
15	H4-1BBRI:	5' CC IAA IGA ACA IGT TTT ACA 3'
		G CT G C G
	H4-1BBRII:	5' TT TTG ATC ATT AAA IGT ICC 3'
20		C G G G

Peripheral blood lymphocytes from normal healthy individuals were isolated and activated with PMA (10 ng/ml) and ionomycin (1 μ M). mRNA from the lymphocytes was isolated. Using reverse transcriptase the human lymphocyte mRNA was converted to single-stranded cDNA. The cDNA was then amplified with Taq polymerase with combination of the primers. The combination of primers was as follows: H4-1BBFI vs H4-1BBRI; H4-1BBFI vs H4-1BBRII; H4-1BBFII vs H4-1BBRI; and H4-1BBFII vs H4-1BBRII.

The primer set of H4-1BBFII and H4-1BBRII produced a specific band of ~240bp. The 240bp is an expected size of human 4-1BB if the human homologue protein is similar to mouse 4-1BB in size. The PCR product (240bp) was cloned in PGEM3 vector and sequenced. One open reading frame of the PCR product was ~65% identical to mouse 4-1BB. Therefore, it was concluded that the 240 bp PCR product is the human homologue of mouse 4-1BB. The 240 bp PCR product was used to screen λ gt11 cDNA library of activated human T lymphocytes. An ~0.85 kb cDNA was isolated. The sequence of the cDNA is shown in Figure 2 and the predicted amino acid sequence is shown in Figure 2b. The same information

is shown is the sequence listing attached to this specification in sequence id. 1.

An expression plasmid to produce H4-1BB-AP fusion
5 protein was constructed. The 5' portion of the H4-1BB cDNA
including sequences encoding the signal sequence and the
entire extracellular domain, was amplified by PCR. For
correctly oriented cloning, a Hind III site on the 5' end
of the forward primer and a Bgl II site on the 5' end of
10 the reverse primer were created.

The Hind III - Bgl II H4-1BB fragment was inserted
into the mammalian expression vector APtaq-1, upstream of
the coding sequence for human placental alkaline
phosphatase (AP). The oligonucleotides PCR primers used
15 for the amplification of 5' portion of H4-1BB are as
follows:

Forward

primer: 5' AAT AAG CTT TGC TAG TAT CAT ACC T 3'

20 Reverse

primer: 5' TTA AGA TCT CTG CGG AGA GTG TCC TGG CTC 3'

H4-1BB-AP will be used to identify cells and tissues
that express ligand for human 4-1BB (i.e. H4-1BBL). The
25 studies with mouse 4-1BB indicated that the ligand for 4-
1BB is on the cell surface. B cells and macrophages were
major cells that express 4-1BBL. It is expected that H4-
1BBL also expresses on human B cells and macrophages.

30 A mammalian expression cDNA library will be generated
from human cell lines that express H4-1BBL. The library
will be screened by [¹²⁵] I-labeled H4-1BB-AP. cDNA for H4-
1BBL will then be isolated and characterized. Soluble
recombinant H4-1BBL will then be produced. Both H4-1BB-AP
35 and H4-1BBL will be used to suppress or enhance immune
responses as described below. Monoclonal antibody to H4-
1BB and H4-1BBL will be produced.

According to studies with mouse 4-1BB, 4-1BB acts as
a costimulatory signal. It is expected that H4-1BB will

act as a costimulatory signal for T cell activation. Mouse 4-1BB helped B cells with proliferation and differentiation. It is expected that H4-1BB will do the same. H4-1BB-AP, H4-1BBL and monoclonal antibody can be
5 used to suppress or enhance human immune responses.

Figures 3a and 3b illustrate the molecules involved in T-cell activation. During early T-cell activation (cognitive phase), resting T cells express the TCR/CD3 complex and other "accessory" molecules. Among these
10 constitutively expressed molecules, CD4 (or CD8), LFA-1 and CD28 are probably the ones to receive costimulatory signals. Initial interaction with the TCR/CD3 complex in combination with these 'accessory' costimulatory signals leads to subsequent expression of additional receptor
15 molecules such as CD28, CTLA4, and 4-1BB. These newly expressed molecules are probably going to receive additional important costimulatory signals at later stages of T-cell activation (clonal expansion).

20 Suppression of immune responses.

Figures 4a-c illustrate a normal T-cell activation pathway. Figures 5a-c illustrate the blocking of immune responses with soluble chimera of 4-1BB. If 4-1BB plays a role in T-cell activation, blocking of the interaction to
25 its ligand on antigen-presenting cells should result in suppression of T-cell dependent immune responses. It is well documented that blocking of the interaction of CD28 to its counter-receptor B7 suppresses in varying degrees, both in vivo antibody production and cell-mediated immune
30 responses. Blocking of both interactions should result in a more effective immunosuppression; since 4-1BB is induced during T-cell activation. Blocking of the interaction of 4-1BB to its ligand may be of importance at later stages of the activation process where the CD28/B7 interaction may no
35 longer be of relevance.

As illustrated with mouse receptor 4-1BB and mouse ligand 4-1BBL above, addition of H4-1BB-AP will coat the H4-

1BBL expressing cells and block the normal interaction between H4-1BB and H4-1BBL. This will lead to immunosuppression. This type of immunosuppression is antigen-specific. Therefore it avoids the generalized immunosuppression produced by antiCD3 or cyclosporin A treatments. H4-1BB-AP treatment can be used to treat certain autoimmune diseases and to facilitate organ transplantation.

10 Immune enhancement.

H4-1BB may function at the late stage of T cell activation and may be a critical molecule for completion of T cell activation. Most tumors display tumor-specific antigens. One reason, however, why immunogenic tumors can escape host immunity is that tumor-reactive T cells receive inadequate costimulation. The introduction of the costimulatory molecules, such as H4-1BB into the tumor, therefore, could enhance the antitumor immunity of cytotoxic T cells (CTL). H4-1BBL can be expressed in cell-specific fashion. For example, the H4-1BBL can be expressed in melanoma using melanocyte-specific promoter such as tyrosinase promoters. The H4-1BBL-expressing melanoma will stimulate cytotoxic T cells through H4-1BB and activate the melanoma-specific CTL. The activated melanoma-specific CTL can destroy melanoma.

Appendix to References Incorporated by Reference

1. Smith, C.A., Davis, T., Anderson, D., Solam, L., Beckmann, M.P., Jerzy, R., Dower, S.K., Cosman, D., and Goodwin, R.G. 1990. A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. Science 248:1019-1023.
2. Ebina, Y., L. Ellis, K. Jaruga, M. Edery, L. Graf, E. Clauser, J. On, F. Marizrz, Y.W. Kan, J.D. Goldfine, R.A. Roth and W.J. Rutter, 1985, The human insulin receptor cDNA: the structural basis for hormone-activated transmembrane signalling, Cell 40:747.
3. Vassali, R., R. Tedghi, B. Listowska-Bernstein, A. Tartakoff and J.C. Jaton, 1979, Evidence for hydrophobic

- region within heavy chains of mouse B lymphocyte membrane-bound IgM, *Proc. Natl. Acad. Sci. USA* 76:5515.
4. Haskins, K., R. Kubo, J. White, M. Pigeon, J. Kappler and P. Marrack, 1983, The major histocompatibility complex-restricted antigen receptor on T cells I Isolation with monoclonal antibody, *J. Exp. Med.* 157:1149.
 5. Lesslaver, W. and H. Gmunder, 1986, Biochemical
10 characterization of the 9.3 antigens of human T-cells: simultaneous expression of disulfide-bonded 90-Kiladalton dimers and free subunits at the cell surface, *Mol. Immunol.* 23:271.
 - 15 6. Van Lier, R., J. Borst, T. Vroom, H. Klein, P. Mourik, W. Zeijlemaker and C. Melife, 1987, Tissue distribution and biochemical and functional properties of Tp55 (CD27) a novel T cell differentiation antigen, *J. Immunol.* 139:1589.
 - 20 7. Mallett, S., S. Fossum and A. Barclay, 1990, Characterization of the MRC OX40 antigen of activated CD4 positive T lymphocytes-a molecule related to nerve growth factor receptor, *EMBO J.* 9:1603.
 - 25 8. Banchereau, J., P. Paoli, A., Valle, E. Garcia and F. Roussel, 1991, Long-term human B cell lines dependent on interleukin-4 and antibody to CD40, *Science* 251:70.
 - 30 9. Moeller, D.L., M.K. Jenkins and R.H. Schwartz, 1989, Clonal expansion versus functional inactivation: a co-stimulatory signalling pathway determines the outcome of T cell antigen receptor occupancy, *Ann. Rev. Immunol.* 7:445.
 - 35 10. June, D.H., J.A. Ledbetter, P.S. Linsley and C.B. Thompson, 1989, Role of CD28 receptor in T cell activation, *Immunol. Today* 11:211.
 - 40 11. Yang, L., B. Jones, A. Aruffo, K.M. Sullivan, P.S. Linsley and C.A. Janeway, Jr., 1992, Heat stable antigen is a co-stimulatory molecule for CD4 T cell growth, *J. Exp. Med.* 175:437.
 - 45 12. Yamori, T., 1992, Molecular mechanisms for generation of neural diversity and specificity: roles of polypeptide factors in development of post-mitotic neurons, *Neurosci. Res.* 12:545.
 - 50 13. Liu, Y.J., D.E. Joshua, G.T. Williams, C.A. Smith, J. Gordon and I.C. M. MacLennan, 1989, Mechanism of antigen-driven selection in germinal centres, *Nature*, 342:929.
 - 55 14. Jabara, H. H., S. M. Fu, R.S. Geha and D. Vercelli, 1990, CD40 and IfE: synergism between anti-CD40 monoclonal antibody and interleukin 4 in the induction of IgE

- synthesis by highly purified human B cells, J. Exp. Med. 172:1861.
15. Defrance, R., B. Vanbervliet, F. Briere, I. Durnad, F. Roussle and J. Banchereau, 1992, Interleukin 10 and transforming growth factor β cooperate to induce anti-CD40 activated naive human B cells to secrete immunoglobulin A, J. Exp. Med. 175:671.
- 10 16. Donahue, T., Cigan, A., Pahich, E. and Valavicius, B., Mutations at a Zn(II) finger motif in the yeast eIF-2 β gene alter ribosomal start-site selection during the scanning process, Cell 54 (1988) 621-632).
- 15 17. Carthew, R.W and Rubin, G.M., seven in absentia, a gene required for specification of R7 cell fate in the Drosophila eye, Cell, 63 (1990) 561-577.
- 20 18. Driscoll, D.M. and Williams, J.G., Two divergently transcribed genes of Dictyostelium discoideum are cyclic AMP-inducible and coregulated during development, Mol. and Cell. Biol. 7 (1987) 4482-4489.
- 25 19. Chalupny, N.J., Peach, R., Hollenbaugh, D., Ledbetter, J.A., Farr, A.G. and Aruffo, A., 1992, Proc. Natl. Acad. Sci USA 89:10360-10364.
- 30 20. Noelle, R.J., and Snow, E.C., 1991, The FASEB J. 5:2770-2776.
21. Noelle, R. and Snow, E., 1990, Immunol. Today 11:361-368.
- 35 22. Zurawski, G., Benedik, M., Kamb, B.J., Abrams, J.S., Zurawski, S.M. and Lee, F.D. (1986) Science 232:772-775.
23. Kinachi, T. (1986) Nature 325,70-73.
- 40 24. Gershenfeld, H.K. and Weissman, I.L. Science (1986) 232,854-858.
25. Biggin, M., Gison, T. and Hung, G. (1983) Proc. Natl. Acad. Sci. USA 80,3963-3965.
- 45 26. Hodgkin, P.D., Yamashita, L.C., Coffman, R.L. and Kehry, M.R., 1990, J. Immunol. 145:2025-2034.
27. Barlett, W.C., McCann, J., Shephaer, D.M., Roy, M. and Noelle, R.J., 1990, J. Immunol. 145:3956-3962.
- 50 28. Kwon, B.S., Kestler, D.P., Eshhar, Z., Oh, K., and Wakulchik, M. 1989. Expression characteristics of two potential T cell mediator genes. Cell. Immunol. 121:414-422.
- 55

29. Armitage, R., Fanslow, W., Strockbine, L., Sato, T., Clifford, K., MacDuff, B., Anderson, D., Gimpel, S., Davis-Smith, T., Maliszewski, C., Clark, E., Smith, C., Grabstein, K., Cosman, D. and Spriggs, M., 191, Nature 5 357:80-82.
30. Kwon, B., Kestler, D., Lee, E., Wakulchik, M. and Young J. (1988) J. Exp. Med (1988) (In press).
- 10 31. Noelle, R.J., Roy, M., Shepherd, D.M., Stamenkovic, I., Ledbetter, J.A. and Aruffo, A., 1992, Proc. Natl. Acad. Sci. USA 89:6550-6554.
- 15 32. Hollenbaugh, D., Grosmaier, L.S., Kullas, C.D., Chalupny, N.J., Braesch-Andersen, S., Noelle, R.J., Stamenkovic, I., Ledbetter, J.A. and Aruffo, A., 1992, EMBO 11:4314-4321.
- 20 33. Schall, T.J., M. Lewis, K.J. Koller, A.Lee, G.C. Rice, G.H. W. Wong, T. Gatanaga, G.A. Granger, R. Lentz, H. Raab, W.J. Kohr and D.V. Goeddel, 1990, Molecular cloning and ex8pression of a receptor for human tumor necrosis factor, Cell 61:361.
- 25 34. Klein, R., Nanduri, V., Jing, S., Lamballe, F., Tapley, P., Bryant, S., Cordon-Cardo, C., Jones, K.R., Reichardt, L.F. and Barbacid, M., 1991, Cell 66:395-403.
- 30 35. Armitage, R.J., Sato, T.A., Macduff, B.M., Clifford, K.N., Alpert, A.R., Smith, C.A. and Fanslow, W.C., 1992, Eu8r. J. Immunol. 22:2071-2076.
- 35 36. Hintzen, R.Q., deJong, R., Hack, E.E., Chamuleau, M., de Vries, E.F.R., ten Berge, I.J.M., Borst, J. and van Lier, R.A.W., 1991, J. Immunol. 147:29-35.
- 40 37. Mallett, S., and Barclay, A.N. 1991. A new super-family of cell surface proteins related to the nerve growth factor receptor. Immunol. Today. 12:220-223.
38. Kwon, B.S., and Weissman, S.M. 1989. cDNA sequences of two inducible T-cell genes. Proc. Natl. Acad. Sci. USA. 86:1963-1967.
- 45 39. Johnson, D., Lanahan, A., Buck C.R., Sehgal, A., Morgan, C., Mercer, E., Bothwell, M., and Chao, M. 1986. Expression and structure of the human NGF receptor. Cell 47:545-554.
- 50 40. Stamenkovic, I., Clark, E., and Seed, B. 1989. A B-lymphocyte activation molecule related to the nerve growth factor receptor and induced by cytokines in carcinomas. EMBO. J. 8:1403-1408.

The foregoing description has been directed to particular embodiments of the invention in accordance with the requirements of the Patent Statutes for the purposes of illustration and explanation. It will be apparent, 5 however, to those skilled in this art that many modifications and changes will be possible without departure from the scope and spirit of the invention. It is intended that the following claims be interpreted to embrace all such modifications.

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
(i) APPLICANT: Byoung Se Kwon
(ii) TITLE OF INVENTION: New Human Receptor and Related Products and Methods
(iii) NUMBER OF SEQUENCES: 1
(iv) CORRESPONDENCE ADDRESS:
 (A) ADDRESSEE: Barnard & Brown
 (B) STREET: 306 E. State St., Suite 220
 (C) CITY: Ithaca
 (D) STATE: New York
 (E) COUNTRY: United States
 (F) ZIP: 14850
(v) COMPUTER READABLE FORM:
 (A) MEDIUM TYPE: Diskette, 3.50 inch, 1.4Mb storage
 (B) COMPUTER: IBM AT Compatible
 (C) OPERATING SYSTEM: MS DOS, Version 5.0
 (D) SOFTWARE: Special QBasic program
(vi) CURRENT APPLICATION DATA:
 (A) APPLICATION NUMBER:
 (B) FILING DATE:
 (C) CLASSIFICATION:
(vii) PRIOR APPLICATION DATA:
 (A) APPLICATION NUMBER: 08/012,269
 (B) FILING DATE: 2/1/93
(viii) PRIOR APPLICATION DATA:
 (A) APPLICATION NUMBER: 07/922,996
 (B) FILING DATE: 7/30/92
(ix) PRIOR APPLICATION DATA:
 (A) APPLICATION NUMBER: 07/267,577
 (B) FILING DATE: 11/7/88
(x) ATTORNEY/AGENT INFORMATION:
 (A) NAME: Michaels, Christopher A.
 (B) REGISTRATION NUMBER: 34,390
 (C) REFERENCE/DOCKET NUMBER: kwnh41bb
(xi) TELECOMMUNICATION INFORMATION:
 (A) TELEPHONE: 607-273-1711
 (B) TELEFAX: 607-273-2609
 (C) TELEX:
(2) INFORMATION FOR SEQ ID NO: 1 :
(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 838
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double stranded
 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA to mRNA
(iii) HYPOTHETICAL: No
(iv) ANTI-SENSE: No
(v) FRAGMENT TYPE: n/a
(vi) ORIGINAL SOURCE:
 (A) ORGANISM: Human
 (B) STRAIN:
 (C) INDIVIDUAL ISOLATE: H4-1BB #1
 (D) DEVELOPMENTAL STAGE: Differentiated T-cell
 (E) HAPLOTYPE:
 (F) TISSUE TYPE:
 (G) CELL TYPE: Lymphocytes
 (H) CELL LINE:
 (I) ORGANELLE:
(vii) IMMEDIATE SOURCE:
 (A) LIBRARY: cDNA library
 (B) CLONE:
(viii) POSITION IN GENOME:
 (A) CHROMOSOME/SEGMENT:
 (B) MAP POSITION:
 (C) UNITS:
(ix) FEATURE:
 (A) NAME/KEY: H4-1BB

(B) LOCATION:
 (C) IDENTIFICATION METHOD: Similarity to mouse 4-1BB and other members of NGFR superfamily
 (D) OTHER INFORMATION:
 (x) PUBLICATION INFORMATION:
 (A) AUTHORS: Kwon, B.S., and Weissman, S.M.
 (B) TITLE: cDNA sequences of two inducible T-cell genes
 (C) JOURNAL: Proc. Natl. Acad. Sci. USA
 (D) VOLUME: 86
 (E) ISSUE:
 (F) PAGES: 1963-1967
 (G) RELEVANT RESIDUES: all

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:
 AATCAGCTTT GCTAGTATCA TACCTGTGCC AGATTTTCATC 40

ATG GGA AAC AGC TGT TAC AAC ATA GTA GCC ACT CTG TTG CTG GTC	85
Met Gly Asn Ser Cys Tyr Asn Ile Val Ala Thr Leu Leu Leu Val	
1 5 10 15	
CTC AAC TTT GAG AGG ACA AGA TCA TTG CAG GAT CCT TGT AGT AAC	130
Leu Asn Phe Glu Arg Thr Arg Ser Leu Gln Asp Pro Cys Ser Asn	
20 25 30	
TGC CCA GCT GGT ACA TTC TGT GAT AAT AAC AGG AAT CAG ATT TGC	175
Cys Pro Ala Gly Thr Phe Cys Asp Asn Asn Arg Asn Gln Ile Cys	
35 40 45	
AGT CCC TGT CCT CCA AAT AGT TTC TCC AGC GCA GGT GGA CAA AGG	220
Ser Pro Cys Pro Pro Asn Ser Phe Ser Ser Ala Gly Gly Gln Arg	
50 55 60	
ACC TGT GAC ATA TGC AGG CAG TGT AAA GGT GTT TTC AGG ACC AGG	265
Thr Cys Asp Ile Cys Arg Gln Cys Lys Gly Val Phe Arg Thr Arg	
65 70 75	
AAG GAG TGT TCC TCC ACC AGC AAT GCA GAG TGT GAC TGC ACT CCA	310
Lys Glu Cys Ser Ser Thr Ser Asn Ala Glu Cys Asp Cys Thr Pro	
80 85 90	
GGG TTT CAC TGC CTG GGG GCA GGA TGC AGC ATG TGT GAA CAG GAT	355
Gly Phe His Cys Leu Gly Ala Gly Cys Ser Met Cys Glu Gln Asp	
95 100 105	
TGT AAA CAA GGT CAA GAA CTG ACA AAA AAA GGT TGT AAA GAC TGT	400
Cys Lys Gln Gly Gln Glu Leu Thr Lys Lys Gly Cys Lys Asp Cys	
110 115 120	
TGC TTT GGG ACA TTT AAC GAT CAG AAA CGT GGC ATC TGT CGA CCC	445
Cys Phe Gly Thr Phe Asn Asp Gln Lys Arg Gly Ile Cys Arg Pro	
125 130 135	
TGG ACA AAC TGT TCT TTG GAT GGA AAG TCT GTG CTT GTG AAT GGG	490
Trp Thr Asn Cys Ser Leu Asp Gly Lys Ser Val Leu Val Asn Gly	
140 145 150	
ACG AAG GAG AGG GAC GTG GTC TGT GGA CCA TCT CCA GCT GAC CTC	535
Thr Lys Glu Arg Asp Val Val Cys Gly Pro Ser Pro Ala Asp Leu	
155 160 165	
TCT CCG GGA GCA TCC TCT GTG ACC CCG CCT GCC CCT GCG AGA GAG	580
Ser Pro Gly Ala Ser Ser Val Thr Pro Pro Ala Pro Ala Arg Glu	
170 175 180	
CCA GGA CAC TCT CCG CAG ATC ATC TCC TTC TTT CTT GCG CTG ACG	625
Pro Gly His Ser Pro Gln Ile Ile Ser Phe Phe Leu Ala Leu Thr	
185 190 195	

TCG	ACT	GCG	TTG	CTC	TTC	CTG	CTG	TTC	TTC	CTC	ACG	CTC	CGT	TTC	670
Ser	Thr	Ala	Leu	Leu	Phe	Leu	Leu	Phe	Phe	Leu	Thr	Leu	Arg	Phe	
				200					205					210	
TCT	GTT	GTT	AAA	CGG	GGC	AGA	AAG	AAA	CTC	CTG	TAT	ATA	TTC	AAA	715
Ser	Val	Val	Lys	Arg	Gly	Arg	Lys	Lys	Leu	Leu	Tyr	Ile	Phe	Lys	
				215					220					225	
CAA	CCA	TTT	ATG	AGA	CCA	GTA	CAA	ACT	ACT	CAA	GAG	GAA	GAT	GGC	760
Gln	Pro	Phe	Met	Arg	Pro	Val	Gln	Thr	Thr	Gln	Glu	Glu	Asp	Gly	
				230					235					240	
TGT	AGC	TGC	CGA	TTT	CCA	GAA	GAA	GAA	GAA	GGA	GGA	TGT	GAA	CTG	805
Cys	Ser	Cys	Arg	Phe	Pro	Glu	Glu	Glu	Glu	Gly	Gly	Cys	Glu	Leu	
				245					250					255	
TGAAATGGAA GTCAATAGGG CTGTTGGGAC TTT															838

□

I claim:

1. A cDNA encoding for human receptor protein H4-1BB.
- 5 2. The cDNA of claim 1 having a nucleotide sequence as shown in Figure 2.
3. The cDNA of claim 1, identified as pH4-1BB deposited at the Agricultural Research Service Culture Collection with
10 the accession number NRRL B21131.
4. The cDNA of claim 2 and fragments and derivatives thereof, wherein said fragments and derivatives can be used as a probe to isolate DNA sequences encoding for proteins
15 similar to the receptor protein encoded by said cDNA.
5. The receptor protein H4-1BB produced by
 - a) inserting the cDNA of H4-1BB into an appropriate expression vector,
 - 20 b) transfecting said expression vector into an appropriate transfection host,
 - c) growing said transfected hosts in appropriate culture media and
 - d) purifying the receptor protein from said culture
25 media.
6. A protein having the amino acid sequence shown in Figure 2.
- 30 7. The protein of claim 6 and fragments and derivatives thereof, wherein said fragments and derivatives:
 - a) can be used as a probe to identify ligands to receptor protein H4-1BB;
 - b) can be used to stimulate proliferation B-cell's
35 expressing H4-1BB ligands; or
 - c) can be used to block H4-1BB ligand binding.

8. A monoclonal antibody against H4-1BB which specifically recognizes receptor protein H4-1BB.
9. A hybridoma capable of producing a monoclonal antibody
5 against H4-1BB which specifically recognizes receptor protein H4-1BB.
10. The method of using the monoclonal antibody of claim 8 to enhance T-cell proliferation comprising the step of
10 treating T-cells that have expressed receptor protein H4-1BB with said monoclonal antibody.
11. The method of claim 12 further comprising the step of
conducting said treatment in the presence of protein
15 tyrosinase kinase.
12. The method of using the monoclonal antibody of claim 8 to enhance T-cell activation comprising the step of
treating T-cells that have expressed receptor protein H4-
20 1BB with said monoclonal antibody.
13. The method of claim 12 further comprising the step of
conducting said treatment in the presence of protein
tyrosinase kinase.
- 25
14. A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB, comprising:
- a) at least a portion of said receptor protein H4-1BB corresponding to the extracellular portion of said
30 receptor protein H4-1BB such that said portion of said receptor protein H4-1BB binds to said cell membrane ligands; and
- b) a detection protein bound to said portion of said receptor protein H4-1BB such that ligand binding can
35 be detected by relative activity assays for said detection protein.

15. The fusion protein of claim 14 wherein said detection protein is alkaline phosphatase.

16. A method of detecting cell membrane ligands to human
5 receptor protein H4-1BB, comprising:

a) providing a fusion protein including:

10 1) at least a portion of said receptor protein H4-1BB corresponding to the extracellular portion of said receptor protein H4-1BB such that said portion of said receptor protein H4-1BB binds to said cell membrane ligands, and

15 2) a detection protein bound to said portion of said receptor protein H4-1BB such that ligand binding can be detected by relative activity assays for said detection protein;

b) placing said fusion protein in the presence of a cell suspected to express said receptor protein H4-1BB;

20 c) washing said cell of any fusion protein not bound to said cell membrane ligands;

d) placing said washed cells in the presence of a substrate for said detection protein and measuring the relative activity of said detection protein.

25 17. The method of claim 16 wherein said detection protein is alkaline phosphatase.

18. A method of inducing B-cell proliferation comprising the step of treating B-cells that have expressed a ligand
30 to human receptor protein H4-1BB with cells that have expressed receptor protein H4-1BB.

-145 CATCAACTGC TGAGTGGATA AACAGGACCG CATACTCTG TCTAAAGCAA TATTACTACA CCAAGGAAAAG ATGTC
 -150 GACACATTGC ACAACAGCAA AGGAGCGTGT CACAGAAAAC CACAGTGTCT TGTGCATGTG ACATTTCGCC
 -70 ATG GGA AAG CAG TGT TAC AAC GTG GTG CTC ATT CTG CTG CTG CTA GTG GCC TGT GAG AAG 60
 1 Met Gly Asn Asn Cys Tyr Asn Val Val Val Ile Val Leu Leu Leu Val Gly Cys Glu Lys 20
 1
 61 GTG GGA GCC GTG CAG AAC TCC TGT GAT AAC TGT CAG CCT GGT ACT TTC TGC AGA AAA TAC 120
 21 Val Gly Ala Val Glu Asn Ser Cys Asp Asn Cys Glu Pro Gly Thr Phe Cys Arg Lys Tyr 40
 121 AAT CCA GTC TGC AAG AGC TGC CCT CCA AGT ACC TTC TCC AGC ATA GGT GGA CAG CCC AAC 180
 41 Asn Pro Val Cys Lys Ser Cys Pro Pro Ser Thr Phe Ser Ser Ile Gly Gly Glu Pro Asn 60
 → H4-IBB F I
 181 TGT AAC ATC TGC AGA GTG TGT GCA CGC TAT TTC AGG TTC AAG AAG TTT TGC TCC TCT ACC 240
 61 Cys Asn Ile Cys Arg Val Cys Ala Gly Tyr Phe Arg Phe Lys Lys Phe Cys Ser Ser Thr 80
 241 CAC AAC GCG GAG TGT CAG TGC ATT GAA GGA TTC CAT TGC TTG GGC CCA CAG TGC ACC AGA 300
 61 His Asn Ala Glu Cys Glu Cys Ile Glu Gly Phe His Cys Leu Gly Pro Glu Cys Thr Arg 100
 301 TGT GAA AAG GAC TGC AGG CCT GGC CAG GAG CTA ACG AAG CAG GGT TGC AAA ACC TGT AGC 360
 101 Cys Glu Lys Asp Cys Arg Pro Gly Glu Glu Leu Thr Lys Glu Gly Cys Lys Thr Cys Ser 120
 361 TTG GGA ACA TTT AAT CAC CAG AAC GGT ACT GGC GTG TGT CGA CCC TGG ACG AAC TCC TCT 420
 121 Leu Gly Thr Phe Asn Asp Glu Asn Gly Thr Gly Val Cys Arg Pro Trp Thr Asn Cys Ser 140
 ← H4-IBB R I
 421 CTA GAC GGA AGC TCT GTG CTT AAG ACC GGC ACC ACG GAG AAG GAC GTG GTG TGT GCA CCC 480
 141 Leu Asp Gly Arg Ser Val Leu Lys Thr Gly Thr Thr Glu Lys Asp Val Val Cys Gly Pro 160
 481 CCT GTG GTG AGC TTC TCT CCC AGT ACC ACC ATT TCT GTG ACT CCA GAG GGA GGA CCA GGA 540
 161 Pro Val Val Ser Phe Ser Pro Ser Thr Thr Ile Ser Val Thr Pro Glu Gly Gly Pro Gly 180
 541 GGG CAC TCC TTG CAG GTC CTT ACC TTG TTC CTG GCG CTG ACA TCG GCT TTG CTG CTG GCC 600
 181 Gly His Ser Leu Glu Val Leu Thr Leu Phe Leu Ala Leu Thr Ser Ala Leu Leu Leu Ala 200
 601 CTG ATC TTC ATT ACT CTC CTG TTC TCT GTG CTC AAA TGG ATC AGG AAA AAA TTC CCC CAC 660
 201 Leu Ile Phe Ile Thr Leu Leu Phe Ser Val Leu Lys Trp Ile Arg Lys Lys Phe Pro His 220
 661 ATA TTC AAG CAA CCA TTT AAG AAG ACC ACT GGA GCA GCT CAA CAG GAA GAT GCT TGT AGC 720
 221 Ile Phe Lys Glu Pro Phe Lys Lys Thr Thr Gly Ala Ala Glu Glu Glu Asp Ala Cys Ser 240
 721 TGC CGA TGT CCA CAG GAA GAA GAA GGA GGA GGA GGC TAT CAG CTG TGA TGTACTATC 780
 241 Cys Arg Cys Pro Glu Glu Glu Glu Gly Gly Gly Gly Gly Tyr Glu Leu ---
 781 CTAGGAGATG TGTGGGCGGA AACCGAGAAG CACTAGGACC CCACCATCCT GTGGAACAGC ACAAGCAACC 850
 851 CCACCACCCT GTCTTACAC ATCATCCTAG ATGATGTGTG GCGCGGCACC TCATCCAACT CTCTTCTAAC 920
 921 GCTAACATAT TTCTCTTTAC CTTTTTAAAT TCCTTTTITA AATTIAAATT TTATGTGTGT GAGTGTTTGT 990
 991 CCTGCCCTGA TGCACAGCTG TGTGTGTGTG TGTGTGTGAC ACTCCTGATG CCTGAGCAGG TCAGAAAGA 1060
 1061 AAGCGTITGGT TCATAAGAA CTGGAGTAT GATGGGCTGT GAGCGCGGnnn GATAGGTCCG GACGGAGACC 1130
 1131 TGTCTTCTTA TTTTAACTG ACTGATATAT AAAAAAAAAA TGATATTTC TGATTTGTAG AGATTTCCT 1200
 1201 GACACCCTTC TAGTTAATGA TCTAAGAGGA ATTGTTGATA CGTAGTATAC TGTATATGTG TATGTATATG 1270
 1271 TATATGTATA TATAAGACT TTTTACTCTC AAAGTCAACC TAGAGTGTCT GTGTACCAGT TCAATTTTAT 1340
 1341 TGCACATTTT ACGTCACACA CACACACACA CACACACACA CACGTTTATA CCGCTACTGT TATCGGTAT 1410
 1411 TCTACCTCAT ATAATGGGAT ACGGTAAAAG GAAACCAAAAG AGTGAGTGAT ATTATTGTGCA CGTGACAGA 1480
 1481 CTACCCCTTC TGGGTACGTA GCGGACAGAC TCCTTCGGCA TGCTTAAAACT TCCCTTAGA AGTCTCGCTCA 1550
 1551 AGTTCGCCGA CGAAGAGCAC AGAGGAGACA CAGTCCGAAA AGTTATTTT TCCGCCAAAT CTTTCCCTGT 1620
 1621 TTGCTGACAC TCCACCCCTT GTGACACATT GAGTGTGATC CTTCGCGCGG AAGGTCAGGT CGTACCCGTC 1690
 1691 TGTAGGGGCG GGGAGACAGA GCGCGCGGGG AGCTACAGAA ATCGACTCAC AGCGCGCCCC GCTATTCCGA 1760
 1761 AATGAACTTA TTTTAACTCT ACAAGTTCG TCCGGGCTCG GCGGACCTAT GCGCTGATC GTTATTACCT 1830
 1831 TATCCTGCGG CCAAGATAAA ACAACCAAAA GCCTTGACTC CGGTACTAAT TCTCCCTGCC GGCGCCGTA 1900
 1901 AGCATAAACG GCGGCTCTCC ACTTTAAGAA CCTTGGCGCG TTCTCGCTGT TCTCGCTTC GTAACCGTT 1970
 1971 CTTACAAAAG TAATTAGTCT TCTTTTCAG CTTCCAAAGT TCTGCTAGT CATCGGACGA TCAAGGCTCG 2040
 2041 TATTTCTTAC GCGTACCCCG TACGCGCCCG CAATAAGCGT ACTGCGCGCG CCGTCAAGG CCGTTTGGTT 2110
 2111 TCAGAACTCC AAGGCGCCCC TCATACCAAC GTTTCGACTT TGATTTCTTC CCGTACGTGC TCGTGGGTGC 2180
 2181 CTTAGATCTT TCTCGTAGT TAT AC

Fig 2a

human homologue of mouse 4-1bb

h4-1bb Length: 838

```

1  AATCAGCTTT GCTAGTATCA TACCIGTGCC AGATTTCATC ATGGGAACA
51  GCCTGTACAA CATAGTAGCC ACCTGTGTGC TGGTCCICAA CTTTGAGAGG
101 ACAAGATCAT TGCAGGATCC TTGTAGTAAC TGCCAGCTG GTACATTCG
151 TGATAATAAC AGGAATCAGA TITGCAGTCC CTGTCCICCA AATAGTTCT
201 CCAGCGCAGG TGCACAAAGG ACCGTGTACA TATGCAGGCA GGTAAAGGT
251 GTTTTCAGGA CCAGGAAGGA GGTTCCTCC ACCAGCAAIG CAGAGTGTA
301 CTGCATCCCA GGTTCCTCT GCTGGGGGC AGGATGCAGC ATGTGTGAAC
351 AGGATTTGTA ACAAGGTCAA GAACTGACAA AAAAAGGTIG TAAAGACTGT
401 TGCYTTGGGA CATTTAACGA TCAGAAACGT GGCATCTGTC GACCTGGAC
451 AAACCTGTTT TTGGAIGGAA AGCTGTGCTT TGTGAATGGG ACCAAGGAGA
501 GGGACGTGGT CTGTGGACCA TCTCCAGCTG ACCCTCTCTC GGGAGCTCC
551 TCTGTGACCC CGCTGCCCC TCGAGAGAGG CCAGGACACT CTCCGCAGAT
601 CATCTCTTTC TTCTTGGCG TACGTCTGAC TCGTTCCTC TTCTGTCTG
651 TCTTCTTCTC GCTCCGTTTC TCTGTGTGTA AACGGGGGAG AAAGAACTC
701 CTGTATATAI TCAACAAACC ATTTATGAGA CCAGTACAAA CTACTAAGA
751 GGAAGATGCC TGTAGCTGCC GATTTCCAGA AGAAGAAGAA GGAGGATGTG
801 AACGTGTAAA TGAAGTCAA TAGGGCTGTT GGGACTTT

```

Fig 2b

```

1  MGNSCYNIVA TLLLVNFER TRSLDPFSN CFAGTCDNN RHOICSPCPP
51  NSYSSAGGOR TCDICROCKG VTRIRKECSS TSAECDCTP GFHCLGAGCS
101 MCEODCKOGO ELTKGCKDC CTGTTHDOKR GICRPUTNCS LDGKSVLVNG
151 TKERDVVCGP SPADLSPGAS SVTPFAPARE PGNSPQIIST FLALTSTALL
201 TLLFLTLRT SVYKRGKKL LYTKQTFHR PVQITQEDG CSCRIFEEEE
251 GGCEL*

```





Fig. 3a




Fig. 3b

4 / 5

NORMAL T-CELL ACTIVATION PATHWAY




Fig. 4a




Fig 4b




Fig. 4c

BLOCKING STEPS IN T-CELL ACTIVATION PATHWAY




Fig. 5a




Fig. 5b




Fig 5c

INTERNATIONAL SEARCH REPORT

Inter. Application No
PCT/US 94/10457

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/12 C12N15/62 C07K14/71 C07K14/715 C07K16/28 G01N33/68		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF IMMUNOLOGY., vol.150, no.3, 1 February 1993, BALTIMORE US pages 771 - 781 POLLOK KE;KIM YJ;ZHOU Z;HURTADO J;KIM KK;PICKARD RT;KWON BS; 'Inducible T cell antigen 4-1BB. Analysis of expression and function.' see the whole document ---	1-18
A	JOURNAL OF IMMUNOLOGY., vol.151, no.3, August 1993, BALTIMORE US pages 1255 - 1262 KIM YJ;POLLOK KE;ZHOU Z;SHAW A;BOHLEN JB;FRASER M;KWON BS; 'Novel T cell antigen 4-1BB associates with the protein tyrosine kinase p56lck1.' see the whole document --- -/--	1-18
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 20 January 1995		Date of mailing of the international search report 15 -02- 1995
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016		Authorized officer Nauche, S

INTERNATIONAL SEARCH REPORT

Inter. nal Application No
PCT/US 94/10457

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol.86, March 1989, WASHINGTON US pages 1963 - 1967 KWON, B.S. ET AL.; 'cDNA sequences of two inducible T-cell genes' see the whole document ---	1-18
P,X	GENE., vol.134, 8 December 1993, AMSTERDAM NL pages 295 - 298 SCHWARZ H;TUCKWELL J;LOTZ M; 'A receptor induced by lymphocyte activation (ILA): a new member of the human nerve-growth-factor/tumor-necrosis-factor receptor family.' see the whole document -----	1-18

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.